

Electron excitation and transfer in photoelectrochemical solar cells

Sørensen, Bent

Publication date:
2001

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Sørensen, B. (2001). *Electron excitation and transfer in photoelectrochemical solar cells*. Poster session presented at Conference on advanced electrochemical systems, Diablerets, Switzerland.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

Electron excitation and transfer in photoelectrochemical solar cells

Bent Sørensen

Physics Department, Roskilde University, DK-4000 Roskilde, Denmark

Email: bes@ruc.dk, Web: <http://mmf.ruc.dk/energy>

In connection with a project aimed at developing industrial production schemes for photoelectrochemical (PEC) solar cells [1], I am trying to improve the understanding of the basic processes involved, by studying the transfer of electrons following absorption in the sensitizer dye, from the central to the peripheral parts of the sensitizer molecule, further on to the surface of the semiconductor electrode, from where it works its way into the regular periodic lattice structure. The electrolyte is believed to contribute to shaping the potential in which the transfer process takes place, both in a bulk way (modelled by treating the electrolyte as a continuous and homogeneous dielectric substance) and through individual ions present in the electrolyte, notably near the semiconductor surface. The electrolyte also mediates the redox process by which the electron transferred to the semiconductor is replaced in the sensitizer dye. This is a slow process likely to limit the overall rate of current formation.

The first step in the quantum mechanical description of the electron transfer is to calculate the excitation of the sensitizer dye caused by the absorption of light. Only this step will be described in the present communication. The example used is an organic coumarin derivative recently synthesised [2]. It has the best absorption properties of any organic sensitizer studied, but falls short of the highest efficiency ruthenium complex dye in use with PEC cells [3].

Figure 1 shows the results of a time-dependent Hartree-Fock calculation of the lower states of the coumarin-derivative [4]. The molecular structure used in the pilot calculation is a basic structure geometrically optimised. However, a Berny optimisation subsequently performed showed modest change in the molecular structure [4]. The Figure shows the difference between the electron densities in the first excited and the ground states. For comparison, the HOMO, LUMO and 2nd lowest excited state molecular orbitals are shown in the right-hand side of the Figure. The orbital gap is a little above 6 eV, while the lowest TDHF excited state is at 1.8 eV, indicating a strongly correlated structure. The Figure shows that the electron density is thinned in the central part of the molecule under excitation, and is regained at the peripheral part, notably around the oxygen atoms of the "arm" added to coumarin-343. This makes the excited electron very easy to transfer to the semiconductor surface. The next excited state is as high as 3.5 eV.


[1] B. Sørensen et al., Production oriented development of photoelectrochemical solar cells, contribution to this conference (2001).

[2] K. Hara et al., A coumarin-derivative dye sensitized nanocrystalline TiO₂ solar cell having a high solar-energy conversion efficiency up to 5.6%. *Chem. Comm.* 569-570, 2001.

[3] S. Zakeeruddin et al., *Inorg. Chemistry*, 36, 5937-5946 (1997).


[4] M. Frisch et al., Gaussian-98 vA.9 software, Gaussian Inc. Pittsburgh 2000.

coumarin343-derivative (basic structure used)
 Electron density (absolute square of wave function) **difference** between 1. excited and ground state, based on TDHF calculation.
 Blue=positive; Purple=negative




$$\Delta E = 1.8 \text{ eV}$$

coumarin343-derivative (basic structure used)
 Electron density (absolute square of wave function) for second lowest unoccupied molecular orbital.
 Colours indicate spin difference.



coumarin343-derivative (basic structure used)
 Electron density (absolute square of wave function) for lowest unoccupied molecular orbital.
 Colours indicate spin difference.



coumarin343-derivative (basic structure used)
 Electron density (absolute square of wave function) for highest occupied molecular orbital.
 Colours indicate spin difference.

